ROSAT HRI observations of Centaurus A

S. Döbereiner
Max-Planck Institut für Extraterrestrische Physik, Garching, Germany

N. Junkes
Astronomisches Institut Potsdam, Germany

S.J. Wagner
Landessternwarte Heidelberg, Germany

H. Zinnecker
Astronomisches Institut Potsdam, Germany

R. Fosbury
Space Telescope - European Coordinating Facility, European Southern Observatory

G. Fabbiano
Harvard-Smithsonian Center for Astrophysics
and

E.J. Schreier
Space Telescope Science Institute

ABSTRACT

We present results from a sensitive high-resolution X-ray observation of the nearby active galaxy Centaurus A (NGC 5128) with the ROSAT HRI. The 65 ksec X-ray image clearly distinguishes different components of the X-ray emission from Cen A: the nucleus and the jet, the diffuse galaxy halo, and a number of individual sources associated with the galaxy. The luminosity of the nucleus increased by a factor of two compared to an earlier ROSAT observation in 1990. The high spatial resolution of the ROSAT HRI shows that most of the knots in the jet are extended both along and perpendicular to the jet axis. We report the detection of a new X-ray feature, at the opposite side of the X-ray jet which is probably due to compression of hot interstellar gas by the expanding southwestern inner radio lobe.

Subject headings: galaxies: active, galaxies: individual (Centaurus A), galaxies: jets, X-rays: galaxies

1. Introduction

The nearby active galaxy Centaurus A (Cen A or NGC 5128) is one of the most interesting objects in the sky. With a distance of 3.5 Mpc (Hui et al. 1993) it is the closest active galaxy to the Milky Way and thus allows us to study different constituents in great detail. The optical morphology is characterized by a more or less normal stellar spheroid, surrounded by a dense ring or disk of dust and gas which has been interpreted as the remains of a recent merger with a smaller dust-rich spiral (e.g. Thomson 1992), which may also induce the nuclear activity. Radio observations reveal an active nucleus and two sets of radio lobes, the outer pair of lobes extending to 9°(several 100 kpc), the inner pair extending to 6′ on the sky. The radio structure of Cen A has been investigated in different spatial scales including the large-scale outer lobes (Junkes et al. 1993), the inner lobes and the jet of the galaxy (Burns, Feigelson, & Schreier 1983, Clarke, Burns, & Norman 1992), and details of the central part at parsec-scale (Tingay 1994). X-ray observations with the EINSTEIN observatory showed a pronounced one-sided X-ray jet, consisting of several knots and extending $\approx 4'$ to the northeast, making it the best-observable X-ray jet of all (Feigelson et al. 1981). With the new sensitive ROSAT HRI observations we are able to examine the X-ray jet and other X-ray emitting components of Cen A in much more detail.

2. Observations

Cen A was observed by ROSAT in three occasions. During the initial calibration phase of ROSAT, Cen A was one of the calibration targets for ROSAT. It was observed for 14159 sec with the ROSAT PSPC and again for 19650 sec with the higher-resolution HRI in July 1990. The quality of these early ROSAT observations was already better than previous observations with the EINSTEIN Observatory. In August 1994 Cen A was reobserved with the ROSAT HRI with an exposure time of 64751 sec.

The ROSAT HRI has a useable field of view of 40′ by 40′ . Over the total field of view, the spatial resolution is on the order of 6″ FWHM. The sensitivity is about 4 times higher than with the EINSTEIN HRI. The useable energy range is 0.1–2.4 keV. The spectral resolution of the ROSAT HRI is nearly nonexistent with $\frac{\Delta E}{E} = 0.7$, but allows to roughly differentiate between hard and soft sources on a relative basis.

3. Components of the X-ray emission of Cen A

The new X-ray data allows us to analyze structures in the central region of Cen A and to compare the X-ray features to radio and optical features, including nucleus, jet, inner radio lobes, the stellar spheroid and the absorbing dust lane.

Fig. 1 shows an enlargement of the X-ray image obtained from the long HRI observation in 1994. The field size is 12.8′ by 12.8′, north is up, east is left. The image is not smoothed to show individual photons.

Fig. 2 shows a overlay of contours from a smoothed X-ray image (smoothed with an adaptive Gaussian filter) on an optical image from the Digitized Palomar Sky Survey.

Fig. 3 is a overlay of the X-ray contours on the 18 cm radio map of Clarke, Burns, & Norman (1992).

In the following we will describe the different X-ray features in comparison to their counterparts in other wavelengths.

3.1. Nucleus and jet

Fig. 4 shows the background-subtracted profile of the jet along the jet axis. The profile was integrated in a box 30″ wide, oriented along the jet (P.A. 55°), using the long observation in Aug. 1994. The background was obtained by extracting profiles in 30″ wide boxes to either side of the jet, averaging, smoothing and scaling them to match the brightness of the underlying diffuse halo southwest of the nucleus.

The nucleus is the bright source at the southwestern end of the jet. The negative residuals after background subtraction southwest of the nucleus are due to the irregular distribution of the absorbing material which shows up in the optical as an irregular dust lane. The profile of knot A may also be affected by this absorption. Judging from the absorption of the diffuse emission near the jet, we estimate the absorption to be of the order of 50 % where the dust lane crosses the jet. The remaining uncertainty in the profile of knot A after background subtraction, due to the irregularity of the dust lane, is estimated to be on the order of 10 % of the peak intensity.

The knot designation follows the nomenclature of Feigelson et al. (1981) and Clarke, Burns, & Norman (1992). In the ROSAT HRI data knots C and D as identified by Feigelson et al. (1981) do not appear as individual knots. These knots also do not show

up in the radio data, whereas all others are clearly discernible (Fig. 3). They were excluded from further analysis. Note, that further interknot X-ray emission can clearly be seen between knots F and G, whereas there is no significant emission outside knot G, where the radio jet blends into the inner northeastern radio lobe.

Fig. 5 shows the transverse profiles of the nucleus and the knots. Superposed on each profile is a template profile obtained from the brightest point source in the field (coincident with a distant early-type galaxy), shifted and scaled approximately to the knots' peak intensities. The template profile is consistent with the HRI point spread function plus an additional artificial extent introduced by the attitude uncertainty of the ROSAT spacecraft (about $2''$ to $3''$ in this observation).

The nucleus of Cen A is compatible with a point source. Knot A is marginally extended in both directions. Knot B is unresolved perpendicular to the jet but clearly extended along the direction of the jet. Knots E, F, and G are diffuse and extended, both along and perpendicular to jet.

Table 1 lists the locations of the nucleus and the knots. The locations correspond to the intensity maxima of the knots. Also listed are the position angles and distances with respect to the nucleus of Cen A.

The X-ray counts of the knots were derived by integrating the background-subtracted profiles shown in Fig. 4. The luminosities were calculated assuming a distance of 3.5 Mpc, a power law spectrum with spectral (energy) index of $\alpha = 0.7$ and an absorbing neutral hydrogen column density of $N_{\rm H} = 10^{21}$ cm^{-2}. Table 1 gives the luminosities in the 0.1 - 2.4 keV spectral range for the nucleus and for the individual knots.

The luminosity in the ROSAT band (0.1-2.4 keV) for the nucleus increased significantly between the shorter observation in July 1990 ($2.7 \pm 1.3 \cdot 10^{38}$ ergs/sec, assuming galactic absorption only) and the longer one in August 1994 ($6.3 \pm 0.6 \cdot 10^{38}$ ergs/sec). This suggests that the bulk of the X-ray emission originates very close to the nucleus. The nucleus is assumed to be strongly absorbed ($N_{\rm H} \approx 10^{23}$cm^{-2}). With such a high absorption, the observable flux would be suppressed by a factor of 500 in the HRI energy range compared to the Galactic foreground absorption, raising the intrinsic nuclear luminosity to more than $\mathrm{L}_X = 2-3 \cdot 10^{41}$ ergs sec^{-1} (0.1-2.4 keV) in 1994. However, this value for $N_{\rm H}$, which is frequently cited in the literature, originates from a spectral fit to an older, spatially unresolved UHURU observation of Cen A (Tucker et al. 1973), and may be substantially in error. ROSAT PSPC spectra of the nucleus are affected by spectral contaminations by emission both of the nearby knot A and of the diffuse halo, and have insufficient counts to constrain the intrinsic $N_{\rm H}$.

The luminosities of the knots within the X-ray jet of Cen A did not change between the 1990 and 1994 observations within the error margins.

The spectral resolution of the HRI allows to detect differences between the amplitude channel spectra of the background and those of the knots and the nucleus. Nucleus and knots are generally harder than the diffuse (background) emission. But it is not sufficient to show significant differences among the spectra of individual knots. The PSPC with its better spectral resolution cannot resolve the individual knots in the X-ray jet. The necessary combination of sufficient spatial and spectral resolution will be provided by future X-ray missions like AXAF or XMM.

3.2. Southwestern X-ray feature

Opposite to the jet, about $5.5'$ southwest of the nucleus, an extended diffuse feature is visible in Fig. 1. The brightest parts show up as an arclike structure just outside of the southwestern inner radio lobe in the contour plot of Fig. 3. The proximity to the outer edge of the radio lobe suggests that the feature consists of hot interstellar matter, probably from the gaseous halo of Cen A, which is compressed or shocked and heated by the expanding inner radio lobe. The polarization map of the inner radio lobes (Fig. 5 of Clarke, Burns, & Norman 1992) shows significant polarization along the outer edge of the southwestern radio lobe, whereas the northeastern radio lobe is less polarized and the polarization direction is not aligned along the outer edge. This supports the idea of interaction between the ISM of Cen A and the southwestern radio lobe (see discussion in Wagner, Döbereiner, & Junkes 1995). A similar feature at the outer boundary of the northeastern lobe cannot be found.

After subtraction of the background and of the bright unrelated point source, the X-ray feature has 660 ± 100 cts in total in the 64751 sec HRI observation. Assuming a distance of 3.5 Mpc, a galactic absorption column $N_{\rm H} = 10^{21}$ cm^{-2}, and a Raymond-Smith spectrum with temperature 0.5 keV and half-

solar metallicity, this corresponds to a total luminosity of $6.4 \pm 1.0 \cdot 10^{38}$ ergs sec^{-1} (0.1 – 2.4 keV).

3.3. Associated point sources

In the total HRI field-of-view we detect about 30 individual point sources with at least 5 σ, using a maximum-likelihood technique. Some of the point sources in the field may be associated with foreground objects. However, the average density of sources in the field is much higher than the source densities observed in similar ROSAT observations away from the galactic plane. Furthermore, the source distribution is centered on Cen A. We estimate that about 70 % to 80 % of them are associated with Cen A. The luminosities of the detected sources are on the order of 10^{38} ergs sec^{-1}. Identification of the sources is in progress.

3.4. Diffuse emission

Figs. 1 and 2 show a roughly circular halo of diffuse emission. It is evident from Fig. 2 that the absorption seen in the X-ray data closely matches the visual absorption indicated by the dust lane. Feigelson et al. (1981) reported the existence of two ridges of diffuse emission parallel to the dust lane, explained by emission from unresolved point sources located in the dust and gas disk. Our data suggest that these ridges are merely the unabsorbed edges of the smooth diffuse halo where they adjoin the absorbed region.

The exact radial profile and total luminosity of the diffuse emission of Cen A has to await a detailed background model for this observation. At this stage it is not possible to differentiate between a hot gaseous halo and integrated emission of unresolved point sources.

However, we can give a rough estimate of $9 \pm 3 \cdot 10^{39}$ ergs sec^{-1} (0.1 - 2.4 keV, assuming 0.5 keV thermal emission and half-solar metallicity) for the total luminosity of the diffuse emission within a radius of 12′, corrected for the absorption by the dust lane. Assuming that all of the diffuse emission comes from a hot gaseous halo, Cen A is still underluminous in this respect compared to the mean (linear) L_X-L_B-relation for early type galaxies (Canizares, Fabbiano, & Trinchieri 1987).

If all the diffuse emission comes from unresolved discrete stellar sources (assuming $\alpha = 0.7$ power law spectra with a different counts-to-flux ratio in comparison to thermal emission-line spectra) the total luminosity in the HRI energy band is calculated to $2 \cdot 10^{40}$ ergs sec^{-1}, which is twice (but compatible within the uncertainties) the luminosity expected for discrete sources within an early-type galaxy of $L_B = 10.5$ (Canizares, Fabbiano & Trinchieri 1987).

4. Summary

With the new X-ray data of Cen A we can improve significantly on previous results an clearly distinguish different components of soft X-ray emission. We give luminosities and profiles for the nucleus and jet components (knots), and report the detection of a new diffuse X-ray feature, probably due to interaction of the southwestern inner radio lobe with the hot ISM of Cen A.

We like to thank the ROSAT team for their outstanding technical and software support.

The ROSAT project is supported by the Bundesministerium für Forschung und Technologie (BMFT/-DARA) and the Max-Planck-Gesellschaft.

The Digitized Sky Survey was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The optical image is based on photographic data obtained with the UK Schmidt Telescope.

REFERENCES

Burns, J. O., Feigelson, E. D., and Schreier, E. J. 1983, ApJ. 273, 128

Canizares, C. R., Fabbiano, G., and Trinchieri, G. 1987, ApJ, 312, 503

Clarke, D. A., Burns, J. O., and Norman, M. L. 1992, ApJ, 395, 444

Feigelson, E. D., Schreier, E. J., Delvaille, J. P., Giacconi, R., Grindlay, J. E., and Lightman, A. P. 1981, ApJ, 251, 31

Hui, X., Ford, H. C., Ciardullo, R., and Jacoby, G. H. 1993, ApJ, 414, 463

Junkes, N., Haynes, R. F., Harnett, J. I., and Jauncey, D. L. 1993, A&A, 269, 29

Thomson, R. C. 1992, MNRAS, 257, 689

Tingay, S. J. 1994, Aust. J. Phys., 47, 619

Tucker, W., Kellogg, E., Gursky, H., Giacconi, R., and Tananbaum, H. 1973, ApJ, 180, 715

Wagner, S. J., Döbereiner, S., and Junkes, N. 1995, in *Röntgenstrahlung from the Universe*, conference proceedings, Würzburg 1995, in press.

This 2-column preprint was prepared with the AAS LaTeX macros v4.0.

Fig. 1.— ROSAT HRI image of Cen A (sum of HRI obervations in 1990 and 1994). The field size is 12.8′ by 12.8′, north is up, east is left. The nucleus is marked with an arrow.

Fig. 2.— X-ray contour map (smoothed with an adaptive Gaussian filter) overlaid on an optical image of Cen A from the Palomar Digitized Sky Survey. The field size is 12.8′ by 12.8′, north is up, east is left.

Fig. 3.— X-ray contour map overlaid on an 18cm radio map (from Clarke, Burns, & Norman 1992). The field size is 12.8′ by 12.8′, north is up, east is left.

Fig. 4.— Background-subtracted profile of the X-ray jet. The nucleus and the brightest knots are marked according to the nomenclature of Feigelson et al. (1987). The bin size along the jet is 1.5″.

Fig. 5.— Background-subtracted transverse profiles of the nuclear source and the knots, offset along the y-axis for clarity. Dashed lines denote the zero lines, the dotted lines represent the profile of a comparison point source scaled to the peak intensity of each source.

TABLE 1

POSITIONS, COUNTRATES, AND LUMINOSITIES OF THE JET COMPONENTS

component or knot	P.A.	dist. (")	integration range[a] (")	total counts	0.1-2.4 keV luminosity (10^{38} ergs/sec)
Nucleus [b]	–	0	-6.75 - 6.75	274 ± 27	6.3 ± 0.6
A	49	14	6.75 - 33.75	236 ± 36	5.4 ± 0.8
B [c]	58	60	33.75 - 93.75	505 ± 80	11.6 ± 1.8
E	56	111	93.75 - 122.25	140 ± 30	3.2 ± 0.7
F	56	136	122.25 - 141.75	69 ± 21	1.6 ± 0.5
between F and G			141.75 - 188.25	76 ± 37	1.8 ± 0.9
G	54	204	188.25 - 210.75	83 ± 15	1.9 ± 0.3
outside G			210.75 - 282.75	26 ± 29	0.6 ± 0.7

[a]Integration in box along P.A. 55^o

[b]Aug. 94 observation. Luminosities calculated under assumption of galactic absorption only

[c]Including locations of knots C and D as observed by EINSTEIN